

Abstract Harmonic Analysis and Group Algebra on the Group AN .

Kahar El-Hussein

*Department of Mathematics, Faculty of Science,
Al Furat University, Dear El Zore, Syria and
Department of Mathematics, Faculty of Arts Science Al Quryyat,
Al-Jouf University, KSA
E-mail : kumath@ju.edu.sa, kumath@hotmail.com*

April 15, 2014

Abstract

Let $G = SL(n, \mathbb{R})$ be the real semisimple Lie group and let $SL(n, \mathbb{R}) = KAN$ be the Iwasawa decomposition. The purpose of this paper is to define the Fourier transform in order to obtain the Plancherel theorem on the group AN . Besides, we prove the existence theorems for the invariant differential operators on the nilpotent group N , and the solvable AN . To this end, we give a classification of all left ideals of the group algebra $L^1(AN)$.

Keywords: Semisimple Lie Groups, Fourier Transform and Plancherel Theorem, Differential Operators, Group Algebra

AMS 2000 Subject Classification: 43A30&35D 05

1 Introduction.

1.1. Abstract harmonic analysis is one of the more modern branches of harmonic analysis, having its roots in the mid-twentieth century, is analysis

on topological groups. The core motivating idea are the various Fourier transforms, which can be generalized to a transform of functions defined on Hausdorff locally compact topological groups. If the group is neither abelian nor compact, no general satisfactory theory is currently known. In mathematics and theoretical physics, the invariant differential operators play very important role in the description of physical symmetries. starting from the early occurrences in the Maxwell, d'Allembert, Dirac, equations. Usually, the action of the group has the meaning of a change of coordinates (change of observer) and the invariance means that the operator has the same expression in all admissible coordinates. For example the Dirac operator in physics is invariant with respect to the Poincaré group and Maxwell equation is invariant by the Lorentz group .

1.2. Thus, it is important for the applications in mathematical physics to study systematically such operators and their solutions. Our interest is to consider the solvable Lie group AN of the real semisimple Lie group $G = SL(n, \mathbb{R})$, Therefore let $SL(n, \mathbb{R}) = KAN$ be the Iwasawa decomposition. In this paper, we will define the Fourier transform on the group AN to obtain the Plancherel formula and to solve any invariant differential operators on the group AN . In the end, we give a classification of all left ideals of group algebra $L^1(AN)$ and we introduce some new groups

2 Notation and Results

2.1. The fine structure of the nilpotent Lie groups will help us to do the Fourier transform on a simply connected nilpotent Lie groups N . As well

known any group connected and simply connected N has the following form

$$N = \begin{pmatrix} 1 & x_1^1 & x_1^2 & x_1^3 & . & . & . & . & . & x_1^{n-2} & x_1^{n-1} & x_1^n \\ 0 & 1 & x_2^2 & x_2^3 & . & . & . & . & . & x_2^{n-2} & x_2^{n-1} & x_2^n \\ 0 & 0 & 1 & x_3^3 & . & . & . & . & . & x_3^{n-2} & x_3^{n-1} & x_3^n \\ 0 & 0 & 0 & 1 & . & . & . & . & . & x_4^{n-2} & x_4^{n-1} & x_4^n \\ . & . & . & . & . & . & . & . & . & . & . & . \\ . & . & . & . & . & . & . & . & . & . & . & . \\ . & . & . & . & . & . & . & . & . & . & . & . \\ . & . & . & . & . & . & . & . & . & . & . & . \\ . & . & . & . & . & . & . & . & . & x_{n-2}^{n-2} & x_{n-2}^{n-1} & x_{n-2}^n \\ 0 & 0 & 0 & 0 & . & . & . & . & . & 1 & x_{n-1}^{n-2} & x_{n-1}^{n-1} \\ 0 & 0 & 0 & 0 & . & . & . & . & . & 0 & 1 & x_n^n \\ 0 & 0 & 0 & 0 & . & . & . & . & . & 0 & 0 & 1 \end{pmatrix} \quad (1)$$

As shown, this matrix is formed by the subgroup $\mathbb{R}, \mathbb{R}^2, \dots, \mathbb{R}^{n-1}$, and $\mathbb{R}^n$

$$\left(\mathbb{R} = \begin{bmatrix} x_1^1 \\ 1 \\ 0 \\ 0 \\ . \\ . \\ . \\ . \\ . \\ 0 \\ 0 \\ 0 \end{bmatrix}, \mathbb{R}^2 = \begin{bmatrix} x_1^2 \\ x_2^2 \\ 1 \\ 0 \\ . \\ . \\ . \\ . \\ . \\ 0 \\ 0 \\ 0 \end{bmatrix}, \dots, \mathbb{R}^{n-1} = \begin{bmatrix} x_1^{n-1} \\ x_2^{n-1} \\ x_3^{n-1} \\ x_4^{n-1} \\ . \\ . \\ . \\ . \\ . \\ x_{n-2}^{n-1} \\ x_{n-2}^{n-2} \\ x_{n-1}^{n-1} \\ 1 \\ 0 \end{bmatrix}, \mathbb{R}^n = \begin{bmatrix} x_1^n \\ x_2^n \\ x_3^n \\ x_4^n \\ . \\ . \\ . \\ . \\ . \\ x_{n-2}^n \\ x_{n-1}^n \\ x_n^n \\ 1 \end{bmatrix} \right) \quad (2)$$

Each $\mathbb{R}^i$ is a subgroup of N of dimension i , $1 \leq i \leq n$, put $d = n + (n - 1) + \dots + 2 + 1 = \frac{n(n+1)}{2}$, which is the dimension of N . According to [9], the group N is isomorphic onto the following group

$$((((\mathbb{R}^n \rtimes_{\rho_n} \mathbb{R}^{n-1}) \rtimes_{\rho_{n-1}} \mathbb{R}^{n-2} \rtimes_{\rho_{n-2}} \dots) \rtimes_{\rho_2} \mathbb{R}^2) \rtimes_{\rho_1} \mathbb{R} \quad (3)$$

That means

$$N \simeq (((((\mathbb{R}^n \rtimes_{\rho_n} \mathbb{R}^{n-1}) \rtimes_{\rho_{n-1}} \mathbb{R}^{n-2} \rtimes_{\rho_{n-2}} \dots) \rtimes_{\rho_4} \mathbb{R}^3 \rtimes_{\rho_3} \mathbb{R}^2) \rtimes_{\rho_2} \mathbb{R} \quad (4)$$

2.2. Denote by $L^1(N)$ the Banach algebra that consists of all complex valued functions on the group N , which are integrable with respect to the Haar measure of N and multiplication is defined by convolution on N as follows:

$$g * f(X) = \int_N f(Y^{-1}X)g(Y)dY \quad (5)$$

for any $f \in L^1(N)$ and $g \in L^1(N)$, where $X = (X^1, X^2, X^3, \dots, X^{n-2}, X^{n-1}, X^n)$, $Y = (Y^1, Y^2, Y^3, \dots, Y^{n-2}, Y^{n-1}, Y^n)$, $X^1 = x_1^1$, $X^2 = (x_1^2, x_2^2)$, $X^3 = (x_1^3, x_2^3, x_3^3)$, $\dots$, $X^{n-2} = (x_1^{n-2}, x_2^{n-2}, x_3^{n-2}, x_4^{n-2}, \dots, x_{n-2}^{n-2})$, $X^{n-1} = (x_1^{n-1}, x_2^{n-1}, x_3^{n-1}, x_4^{n-1}, \dots, x_{n-2}^{n-1}, x_{n-1}^{n-1})$, $X^n = (x_1^n, x_2^n, x_3^n, x_4^n, \dots, x_{n-2}^n, x_{n-1}^n, x_n^n)$ and $dY = dY^1 dY^2 dY^3, \dots, dY^{n-2} dY^{n-1} dY^n$ is the Haar measure on N and $*$ denotes the convolution product on N . We denote by $L^2(N)$ its Hilbert space. Let $\mathcal{U}$ be the complexified universal enveloping algebra of the real Lie algebra $\underline{g}$ of N ; which is canonically isomorphic to the algebra of all distributions on N supported by $\{0\}$, where 0 is the identity element of N . For any $u \in \mathcal{U}$ one can define a differential operator P_u on N as follows:

$$P_u f(X) = u * f(X) = \int_N f(Y^{-1}X)u(Y)dY \quad (6)$$

The mapping $u \rightarrow P_u$ is an algebra isomorphism of $\mathcal{U}$ onto the algebra of all invariant differential operators on N

3 Fourier Analysis on N .

3.1. For each $2 \leq i \leq n$, let

$$K_{i-1} = \left\{ \left\{ \underbrace{\left\{ \underbrace{\mathbb{R}^n}_{\rtimes_{\rho_n}} \times \mathbb{R}^{n-1} \times \mathbb{R}^{n-1} \right\} \times \mathbb{R}^{n-2} \times \mathbb{R}^{n-2} \times \dots}_{\rtimes_{\rho_{n-1}}} \times \mathbb{R}^i \times \mathbb{R}^i \right\} \times \mathbb{R}^{i-1} \rtimes_{\rho_i} \mathbb{R}^{i-1} \right\}$$

be the group with the following law

$$\begin{aligned} & (X, \dots, X^i, X^i, U^{i-1}, X^{i-1}).(Y, \dots, Y^i, Y^i, V^{i-1}, Y^{i-1}) \\ &= [(X, \dots, X^i, X^i).(\rho_i(X^i)(Y, \dots, Y^i, Y^i))], U^{i-1} + V^{i-1}, X^{i-1} + Y^{i-1} \end{aligned} \quad (7)$$

For any $X \in L_{i+1}, Y \in L_{i+1}, X^i \in \mathbb{R}^i, U^{i-1} \in \mathbb{R}^{i-1}, X^{i-1} \in \mathbb{R}^{i-1}, Y^i \in \mathbb{R}^i, V^{i-1} \in \mathbb{R}^{i-1}, Y^{i-1} \in \mathbb{R}^{i-1}$. So we get for $i = 2$

$$K_1 = \left\{ \left\{ \underbrace{\left\{ \underbrace{\mathbb{R}^n}_{\rtimes_{\rho_n}} \times \mathbb{R}^{n-1,1} \rtimes_{\rho_{n-1}} \mathbb{R}^{n-1} \right\} \times \mathbb{R}^{n-2,1} \rtimes_{\rho_3} \mathbb{R}^{n-2} \times \dots}_{\rtimes_{\rho_{n-1}}} \times \mathbb{R}^{2,1} \times \mathbb{R}^2 \right\} \times \mathbb{R}^{1,1} \rtimes_{\rho_2} \mathbb{R}^1 \right\}$$

and for $i = n$, we get

$$K_{n-1} = \left\{ \underbrace{\mathbb{R}^n}_{\rtimes_{\rho_n}} \right\} \times \mathbb{R}^{n-1} \rtimes_{\rho_n} \mathbb{R}^{n-1}$$

3.2. Let $M = \mathbb{R}^n \times \mathbb{R}^{n-1,1} \times \mathbb{R}^{n-2,1} \times \dots \times \mathbb{R}^{3,1} \times \mathbb{R}^{2,1} \times \mathbb{R}^{1,1} = \mathbb{R}^d$ be the Lie group, which is the direct product of $\mathbb{R}^{n,1} = \mathbb{R}^n$, $\mathbb{R}^{n-1,1} = \mathbb{R}^{n-1}$, $\mathbb{R}^{n-2,1} = \mathbb{R}^{n-2}$, $\dots$, $\mathbb{R}^{3,1} = \mathbb{R}^3$, $\mathbb{R}^{2,1} = \mathbb{R}^2$ and $\mathbb{R}^{1,1} = \mathbb{R}^1$. Denote by $L^1(M)$ the Banach algebra that consists of all complex valued functions on the group M , which are integrable with respect to the Lebesgue measure on M and multiplication is defined by convolution on M as:

$$g *_c f(X) = \int_M f(X - Y)g(Y)dY \quad (8)$$

for any $f \in L^1(M)$, $g \in L^1(M)$, where $*_c$ signifies the convolution product on the abelian group M . We denote again by $\mathcal{U}$ be the complexified universal enveloping algebra of the real Lie algebra $\underline{m}$ of M ; which is canonically isomorphic to the algebra of all distributions on M supported by $\{0\}$, where 0 is the identity element of M . For any $u \in \mathcal{U}$ one can define a differential operator Q_u on M as follows:

$$Q_u f(X) = u *_c f(X) = f *_c u(X) = \int_M f(X - Y)u(Y)dY \quad (9)$$

The mapping $u \rightarrow Q_u$ is an algebra isomorphism of $\mathcal{U}$ onto the algebra of all invariant differential operators (the algebra of all linear differential operators with constant coefficients) on M

The group N can be identified with the subgroup

$$N = \left\{ \underbrace{\left\{ \underbrace{\left\{ \underbrace{\mathbb{R}^n}_{\rtimes_{\rho_n}} \right\} \times \{0\}}_{\rtimes_{\rho_{n-1}}} \rtimes_{\rho_{n-1}} \mathbb{R}^{n-1} \right\} \times \{0\} \rtimes_{\rho_3} \mathbb{R}^{n-2} \times \dots}_{\rtimes_{\rho_2}} \right\} \times \{0\} \rtimes_{\rho_2} \mathbb{R} \quad (10)$$

of K_1 and M can be identified with the subgroup

$$M = \left\{ \underbrace{\left\{ \underbrace{\left\{ \underbrace{\mathbb{R}^n}_{\rtimes_{\rho_n}} \right\} \times \mathbb{R}^{n-1,1} \rtimes_{\rho_n} \{0\}}_{\rtimes_{\rho_{n-1}}} \right\} \times \mathbb{R}^{n-2,1} \rtimes_{\rho_{n-1}} \{0\} \times \dots}_{\rtimes_{\rho_2}} \right\} \times \mathbb{R}^{1,1} \rtimes_{\rho_2} \{0\}$$

In this paper, we show how the Fourier transform on the vector group $\mathbb{R}^d$ can be generalized on N and obtain the Plancherel theorem.

Definition 3.1. For $1 \leq i \leq n$, let $\mathcal{F}^i$ be the Fourier transform on $\mathbb{R}^i$ and $0 \leq j \leq n-1$, let $\prod_{0 \leq l \leq j} \mathbb{R}^{n-l} = (..(((\mathbb{R}^n \rtimes_{\rho_n} \mathbb{R}^{n-1}) \rtimes_{\rho_{n-1}}) \mathbb{R}^{n-2} \rtimes_{\rho_{n-2}}) .. \rtimes_{\rho_{n-j}} \mathbb{R}^{n-j})$, and let $\prod_{0 \leq l \leq j} \mathcal{F}^{n-l} = \mathcal{F}^n \mathcal{F}^{n-1} \mathcal{F}^{n-2} \dots \mathcal{F}^{n-j}$, we can define the Fourier

$$\begin{aligned}
& \text{transform on } \prod_{0 \leq l \leq n-1} \mathbb{R}^{n-1} = \mathbb{R}^n \rtimes_{\rho_n} \mathbb{R}^{n-1} \rtimes_{\rho_{n-1}} \mathbb{R}^{n-2} \rtimes_{\rho_{n-2}} \dots \rtimes_{\rho_3} \mathbb{R}^2 \rtimes_{\rho_2} \mathbb{R}^1 \text{ as} \\
& \mathcal{F}^n \mathcal{F}^{n-1} \mathcal{F}^{n-2} \dots \mathcal{F}^2 \mathcal{F}^1 f(\lambda^n, \lambda^{n-1}, \lambda^{n-2}, \dots, \lambda^2, \lambda^1) \\
& = \int_N f(X^n, X^{n-1}, \dots, X^2, X^1) e^{-i \langle (\lambda^n, \lambda^{n-1}), (X^n, X^{n-1}), \dots, (\lambda^2, \lambda^1), (X^2, X^1) \rangle} \\
& \quad dX^n dX^{n-1} \dots dX^2 dX^1 \tag{11}
\end{aligned}$$

for any $f \in L^1(N)$, where $X = (X^n, X^{n-1}, \dots, X^2, X^1)$, $\mathcal{F}^d = \mathcal{F}^n \mathcal{F}^{n-1} \mathcal{F}^{n-2} \dots \mathcal{F}^2 \mathcal{F}^1$ is the classical Fourier transform on N , $dX = dX^n dX^{n-1} \dots dX^2 dX^1$, $\lambda = (\lambda^n, \lambda^{n-1}, \lambda^{n-2}, \dots, \lambda^2, \lambda^1)$, and

$$\begin{aligned}
& \langle (\lambda^n, \lambda^{n-1}), (X^n, X^{n-1}), \dots, (\lambda^2, \lambda^1), (X^2, X^1) \rangle = \sum_{i=1}^n X_i^n \lambda_i^n + \sum_{j=1}^{n-1} X_j^{n-1} \lambda_j^{n-1} + \\
& \dots + \sum_{i=1}^2 X_i^2 \lambda_i^2 + X^1 \lambda^1
\end{aligned}$$

Definition 3.2. For every function f on N one can define a function $\tilde{f}$ on K_1 as

$$\tilde{f}(\rho_2(X^1)[(\rho_3(X^2)(\dots(\rho_{n-1}(X^{n-2})((\rho_n(X^{n-1})(X^n))), U^{i-1} + X^{i-1}), U^{i-2} + X^{i-2}), \dots), U^2 + X^2], U^1 + X^1)$$

Remark 3.1. The function $\tilde{f}$ is invariant in the following sense

$$\begin{aligned}
& \tilde{f}(\rho_2(U^1)[(\rho_3(U^2)(\dots((\rho_n(U^{n-1})(X^n)), X^{n-1} - U^{n-1}), X^{n-2} - U^{n-2}), \dots), X^2 - U^2], X^1 - U^1) \\
& = \tilde{f}(X, U^{i-1}, X^{i-1}, U^{i-2}, X^{i-2}, \dots, U^2, X^2, U^1, X^1) \tag{12}
\end{aligned}$$

Theorem 3.1. For every function F on K_1 invariant in sense and for every function ϕ on N

$$\begin{aligned}
& \phi * F(X, U^{n-1}, X^{n-1}, U^{n-2}, X^{n-2}, \dots, U^2, X^2, U^1, X^1) \\
& = \phi *_c F(X, U^{n-1}, X^{n-1}, U^{n-2}, X^{n-2}, \dots, U^2, X^2, U^1, X^1) \tag{13}
\end{aligned}$$

for every $(X, U^{n-1}, X^{n-1}, U^{n-2}, X^{n-2}, \dots, U^2, X^2, U^1, X^1) \in K_1$, where $*$ signifies the convolution product on N with respect the variables $(X, X^{i-1}, X^{i-2}, \dots, X^2, X^1)$ and $*_c$ signifies the commutative convolution product on M with respect the variables $(X, U^{i-1}, U^{i-2}, \dots, U^2, U^1)$

Proof:

$$\begin{aligned}
& \phi * F(X, U^{n-1}, X^{n-1}, U^{n-2}, X^{n-2}, \dots, U^2, X^2, U^1, X^1) \\
= & \int_N F[(Y^n, Y^{n-1}, Y^{n-2}, \dots, Y^2, Y^1)^{-1}(X, U^{n-1}, X^{n-1}, U^{n-2}, X^{n-2}, \dots, U^2, X^2, U^1, X^1)] \\
& \phi(Y^n, Y^{n-1}, Y^{n-2}, \dots, Y^2, Y^1) dY dY^{i-1} dY^{i-2} \dots dY^2 dY^1 \\
= & \int_N F((\rho(-Y^1)(Y^n, Y^{n-1}, Y^{n-2}, \dots, Y^2)^{-1}), -Y^1)(X, U^{n-1}, X^{n-1}, U^{n-2}, X^{n-2}, \dots, U^2, X^2, U^1, X^1) \\
& \phi(Y, Y^n, Y^{n-1}, Y^{n-2}, \dots, Y^2, Y^1) dY dY^{n-1} dY^{n-2} \dots dY^2 dY^1 \\
= & \int_N F((\rho(-Y^1)((Y^n, Y^{n-1}, Y^{n-2}, \dots, Y^2)^{-1}(X, U^{n-1}, X^{n-1}, U^{n-2}, X^{n-2}, \dots, U^2, X^2, U^1, X^1)), X^1 - \\
& \phi(Y^n, Y^{n-1}, Y^{n-2}, \dots, Y^2, Y^1) dY^n dY^{n-1} dY^{n-2} \dots dY^2 dY^1 \\
= & \int_N F(\rho_2(-Y^1)(\rho_3(-Y^2)((Y^n, Y^{n-1}, Y^{n-2}, \dots, Y^2)^{-1}(X^n, U^{n-1}, X^{n-1}, U^{n-2}, X^{n-2}, \dots, U^2, X^2 - Y^2, \\
& U^1, X^1 - Y^1)] \phi(Y^n, Y^{n-1}, Y^{n-2}, \dots, Y^2, Y^1) dY^n dY^{n-1} dY^{n-2} \dots dY^2 dY^1
\end{aligned}$$

By a simple calculation, we find

$$\begin{aligned}
& \phi * F(X^n, U^{i-1}, X^{i-1}, U^{i-2}, X^{i-2}, \dots, U^2, X^2, U^1, X^1) \\
= & \int_N F[(X^n - Y^n, U^{n-1}, X^{n-1} - Y^{n-1}, U^{n-2}, X^{n-2} - Y^{n-2}, \dots, U^2, X^2 - Y^2, U^1, X^1 - Y^1)] \\
& \phi(Y^n, Y^{i-1}, Y^{i-2}, \dots, Y^2, Y^1) dY^n dY^{i-1} dY^{i-2} \dots dY^2 dY^1 \\
= & \int_M F[(X^n - Y^n, U^{n-1} - Y^{n-1}, X^{n-1}, U^{n-2} - Y^{n-2}, X^{n-2}, \dots, U^2 - Y^2, X^2, U^1 - Y^1, X^1)] \\
& \phi(Y^n, Y^{n-1}, Y^{n-2}, \dots, Y^2, Y^1) dY^n dY^{n-1} dY^{n-2} \dots dY^2 dY^1 \\
= & \phi *_c F(X^n, U^{n-1}, X^{n-1}, U^{n-2}, X^{n-2}, \dots, U^2, X^2, U^1, X^1) \tag{14}
\end{aligned}$$

Corollary. For every function $F \in L^1(K_1)$ invariant in sense and for every function $\phi \in L^1(N)$

$$\begin{aligned}
& \int \int \dots \int \int \mathcal{F}^d(\phi * F)(\xi^n, \lambda^{n-1}, \mu^{n-1}, \lambda^{n-2}, \mu^{n-2}, \dots, \lambda^2, \mu^2, \lambda^1, \mu^1) d\mu^{n-1} d\mu^{n-2} \dots d\mu^2 d\mu^1 \\
= & \mathcal{F}^d F(\xi^n, \lambda^{n-1}, 0, \lambda^{n-2}, 0, \dots, \lambda^2, 0, \lambda^1, 0) \mathcal{F}^d \phi(\xi^n, \lambda^{n-1}, \lambda^{n-2}, \dots, \lambda^2, \lambda^1) \tag{15}
\end{aligned}$$

Proof: In fact we get

$$\begin{aligned}
& \int \int \dots \int \int \mathcal{F}^d(\phi * F)(\xi^n, \lambda^{n-1}, \mu^{n-1}, \lambda^{n-2}, \mu^{n-2}, \dots, \lambda^2, \mu^2, \lambda^1, \mu^1) d\mu^{n-1} d\mu^{n-2} \dots d\mu^2 d\mu^1 \\
= & \int \int \dots \int \int \mathcal{F}^d(\phi *_c F)(\xi^n, \lambda^{n-1}, \mu^{n-1}, \lambda^{n-2}, \mu^{n-2}, \dots, \lambda^2, \mu^2, \lambda^1, \mu^1) d\mu^{n-1} d\mu^{n-2} \dots d\mu^2 d\mu^1 \\
= & \int \int \dots \int \int \mathcal{F}^d F(\xi^n, \lambda^{n-1}, \mu^{n-1}, \lambda^{n-2}, \mu^{n-2}, \dots, \lambda^2, \mu^2, \lambda^1, \mu^1) \mathcal{F}^d \phi(\xi^n, \lambda^{n-1}, \lambda^{n-2}, \dots, \lambda^2, \lambda^1) \\
& d\mu^{n-1} d\mu^{n-2} \dots d\mu^2 d\mu^1 \\
= & \mathcal{F}^d F(\xi^n, \lambda^{n-1}, 0, \lambda^{n-2}, 0, \dots, \lambda^2, 0, \lambda^1, 0) \mathcal{F}^d \phi(\xi^n, \lambda^{n-1}, \lambda^{n-2}, \dots, \lambda^2, \lambda^1)
\end{aligned}$$

Plancherels theorem 3.2. For any function $f \in L^1(N)$, we have

$$\begin{aligned}
& \int_N |\mathcal{F}^d f(\xi^n, X^{n-1}, X^{n-2}, \dots, X^2, X^1)|^2 dX dX^{n-1} dX^{n-2} \dots dX^2 dX^1 \\
& \int_N |\mathcal{F}^d f(\xi^n, \lambda^{n-1}, \lambda^{n-2}, \dots, \lambda^2, \lambda^1)|^2 d\xi^n d\lambda^{n-1} d\lambda^{n-2} \dots d\lambda^2 d\lambda^1 \quad (16)
\end{aligned}$$

Proof: Let $f \in L^1(K_1)$ be the function defined as

$$\begin{aligned}
& \widetilde{\vee} f(X^n, U^{n-1}, X^{n-1}, U^{n-2}, X^{n-2}, \dots, U^2, X^2, U^1, X^1) \\
= & \vee f[(\rho_2(X^1)((\rho_3(X^2)(\dots((\rho_n(X^{n-1})(X^n)), U^{n-1} + X^{n-1}), U^{n-2} + X^{n-2}) \dots), U^2 + X^2)), U^1 + X^1) \\
= & f[(\rho_2(X^1)((\rho_3(X^2)(\dots((\rho_n(X^{n-1})(X^n)), U^{n-1} + X^{n-1}), U^{n-2} + X^{n-2}) \dots), U^2 + X^2)), U^1 + X^1)
\end{aligned}$$

then we have

$$\begin{aligned}
& \phi * \widetilde{\phi}(0, 0, 0, 0, 0, ..0, 0, 0, 0) \\
= & \int_N \widetilde{\phi} [(X^n, X^{i-1}, X^{i-2}, ..., X^2, X^1)^{-1}(0, 0, 0, 0, 0, ..0, 0, 0, 0)] \phi(X^n, X^{i-1}, X^{i-2}, ..., X^2, X^1) \\
& dX^n dX^{n-1} dX^{n-2} ..dX^2 dX^1 \\
= & \int_N \widetilde{\phi} (\rho_2(X^1)[(\rho_3(X^2)(..(\rho_{n-1}(X^{n-2})((\rho_n(X^{n-1})(-X^n)), 0, -X^{i-1}), 0, -X^{i-2}, ...,)0, -X^2,) \\
& 0, -X^1)]\phi(X^n, X^{i-1}, X^{i-2}, ..., X^2, X^1) dX^n dX^{n-1} dX^{n-2} ..dX^2 dX^1 \\
= & \int_N \widetilde{\phi} [(\rho_2(X^1)[(\rho_3(X^2)(..(\rho_{n-1}(X^{n-2})((\rho_n(X^{n-1})(-X^n)), -X^{i-1}), -X^{i-2}, ...,), -X^2,), -X^1)] \\
& \phi(X^n, X^{i-1}, X^{i-2}, ..., X^2, X^1) dX^n dX^{n-1} dX^{n-2} ..dX^2 dX^1 \\
= & \int_N \widetilde{\phi} [(X^n, X^{i-1}, X^{i-2}, ..., X^2, X^1)^{-1}] \phi(X^n, X^{i-1}, X^{i-2}, ..., X^2, X^1) dX^n dX^{n-1} dX^{n-2} ..dX^2 dX^1 \\
= & \int_N \overline{\phi(X^n, X^{i-1}, X^{i-2}, ..., X^2, X^1)} \phi(X^n, X^{i-1}, X^{i-2}, ..., X^2, X^1) dX^n dX^{n-1} dX^{n-2} ..dX^2 dX^1 \\
= & \int_N |\phi(X^n, X^{i-1}, X^{i-2}, ..., X^2, X^1)|^2 dX^n dX^{n-1} dX^{n-2} ..dX^2 dX^1
\end{aligned}$$

In other hand we have

$$\begin{aligned}
& \phi * \widetilde{\phi}(0, 0, 0, 0, 0, 0, \dots, 0, 0, 0, 0) \\
&= \int \mathcal{F}^d(\phi * \widetilde{\phi})(\xi^n, \lambda^{n-1}, \mu^{n-1}, \lambda^{n-2}, \mu^{n-2}, \dots, \lambda^2, \mu^2, \lambda^1, \mu^1) \\
&\quad d\xi^n d\lambda^{n-1} d\mu^{n-1} d\lambda^{n-2} d\mu^{n-2} \dots d\lambda^2 d\mu^2 d\lambda^1 d\mu^1 \\
&= \int \mathcal{F}^d(\phi * \widetilde{\phi})(\xi^n, \lambda^{n-1}, 0, \lambda^{n-2}, 0, \dots, \lambda^2, 0, \lambda^1, 0) d\xi^n d\lambda^{n-1} d\lambda^{n-2} \dots d\lambda^2 d\lambda^1 \\
&= \int \mathcal{F}^d(\phi * \widetilde{\phi})(\xi^n, \lambda^{n-1}, 0, \lambda^{n-2}, 0, \dots, \lambda^2, 0, \lambda^1, 0) d\xi^n d\lambda^{n-1} d\lambda^{n-2} \dots d\lambda^2 d\lambda^1 \\
&= \int \mathcal{F}^d \widetilde{\phi}(\xi^n, \lambda^{n-1}, 0, \lambda^{n-2}, 0, \dots, \lambda^2, 0, \lambda^1, 0) \mathcal{F}^d \phi(\xi^n, \lambda^{n-1}, \lambda^{n-2}, \dots, \lambda^2, \lambda^1) \\
&\quad d\xi^n d\lambda^{n-1} d\lambda^{n-2} \dots d\lambda^2 d\lambda^1 \\
&= \int \overline{\mathcal{F}^d \phi(\xi^n, \lambda^{n-1}, \lambda^{n-2}, \dots, \lambda^2, \lambda^1)} \mathcal{F}^d \phi(\xi^n, \lambda^{n-1}, \lambda^{n-2}, \dots, \lambda^2, \lambda^1) d\xi^n d\lambda^{n-1} d\lambda^{n-2} \dots d\lambda^2 d\lambda^1 \\
&= \int |\mathcal{F}^d \phi(\xi^n, \lambda^{n-1}, \lambda^{n-2}, \dots, \lambda^2, \lambda^1)|^2 d\xi^n d\lambda^{n-1} d\lambda^{n-2} \dots d\lambda^2 d\lambda^1
\end{aligned}$$

Hence the proof of the theorem.

3.1 Division of Distributions on N .

3.1.1. Let $\mathcal{S}(N)$ be the Schwartz space of N which can be considered as the Schwartz space of $\mathcal{S}(M)$, let $\mathcal{S}'(N)$ be the space of all tempered distributions on N and let $\mathcal{S}'(M)$ be the space of all tempered distributions on M . If we consider the group N as a subgroup of K_1 , then $\tilde{f} \in \mathcal{S}(N)$ for $U = (U^{n-1}, U^{n-2}, \dots, U^2, U^1)$ is fixed, and if we consider M as a subgroup of K_1 , then $\tilde{f} \in \mathcal{S}(M)$ for $X = (X^{n-1}, X^{n-2}, \dots, X^2, X^1)$ is fixed. This being so; denote by $\mathcal{S}_E(K_1)$ the space of all functions $\Phi(X^n, U^{n-1}, X^{n-1}, U^{n-2}, X^{n-2}, \dots, U^2, X^2, U^1, X^1) \in C^\infty(K_1)$ such that $\Phi(X^n, U^{n-1}, X^{n-1}, U^{n-2}, X^{n-2}, \dots, U^2, X^2, U^1, X^1) \in \mathcal{S}(N)$ for $X = (X^{n-1}, X^{n-2}, \dots, X^2, X^1)$ is fixed, and $\Phi(X^n, U^{n-1}, X^{n-1}, U^{n-2}, X^{n-2}, \dots, U^2, X^2, U^1, X^1) \in \mathcal{S}(M)$ for $U = (U^{n-1}, U^{n-2}, \dots, U^2, U^1)$ is fixed. We equip $\mathcal{S}_E(K_1)$ with the natural topology defined by the seminorms:

$$\Phi \rightarrow \sup_{(X^n, U) \in M} |Q(Y)P(D)\Phi(X^n, U^{n-1}, X^{n-1}, U^{n-2}, X^{n-2}, \dots, U^2, X^2, U^1, X^1)| \quad X \text{ fixed}$$

$$\Phi \rightarrow \sup_{(X^n, X) \in N} |R(Y)S(D)\Phi(X^n, U^{n-1}, X^{n-1}, U^{n-2}, X^{n-2}, \dots, U^2, X^2, U^1, X^1)| \quad U \text{ fixed}$$

where $(X^n, X) = (X^n, X^{n-1}, X^{n-2}, \dots, X^2, X^1)$, $(X^n, U) = (X^n, U^{n-1}, U^{n-2}, \dots, U^2, U^1)$, and P, Q, R and S run over the family of all complex polynomials in $2d - n$ variables. Let $\mathcal{S}_E^I(K_1)$ be the subspace of all functions $\psi \in \mathcal{S}_E(K_1)$, which are invariant in sense (13), then we have the following result.

Lemma 3.1.1. *Let $u \in \mathcal{U}$ and Q_u be the invariant differential operator on the group M , which is associated to u , acts on the variables $(X^n, U^{n-1}, U^{n-2}, \dots, U^2, U^1) \in M$, then we have*

(i) *The mapping $f \mapsto \tilde{f}$ is a topological isomorphism of $\mathcal{S}(N)$ onto $\mathcal{S}_E^I(K_1)$.*

(ii) *The mapping $\Phi \mapsto Q_u \Phi$ is a topological isomorphism of $\mathcal{S}_E^I(K_1)$ onto its image*

Proof: (i) In fact $\sim$ is continuous and the restriction mapping $\Phi \mapsto R\Phi$ on N is continuous from $\mathcal{S}_E^I(K_1)$ into $\mathcal{S}(N)$ that satisfies $R \circ \sim = Id_{\mathcal{S}(N)}$ and $\sim \circ R = Id_{\mathcal{S}_E^I(K_1)}$, where $Id_{\mathcal{S}(N)}$ (resp. $Id_{\mathcal{S}_E^I(K_1)}$) is the identity mapping of $\mathcal{S}(N)$ (resp. $\mathcal{S}_E^I(K_1)$) and N is considered as a subgroup of K_1 .

To prove(ii) we refer to [16, P.313 – 315] and his famous result that is: "Any invariant differential operator on M , is a topological isomorphism of $\mathcal{S}(M)$ onto its image" From this result, we obtain that

$$D_u : \mathcal{S}_E(K_1) \rightarrow \mathcal{S}_E(K_1) \quad (17)$$

is a topological isomorphism and its restriction on $\mathcal{S}_E^I(K_1)$ is a topological isomorphism of $\mathcal{S}_E^I(K_1)$ onto its image. Hence the theorem is proved.

In the following we will prove that every invariant differential operator on N has a tempered fundamental solution. As in the introduction, we will consider the two invariant differential operators P_u and Q_u , the first on the group N , and the second on the abelian vector group M . Our main result is:

Theorem 3.1.1. *Every nonzero invariant differential operator on N has a tempered fundamental solution*

Proof: For every function $\psi \in C^\infty(K_1)$ invariant in sense (11) and for

every $u \in \mathcal{U}$, we have

$$\begin{aligned}
& P_u \psi(X^n, U^{n-1}, X^{n-1}, U^{n-2}, X^{n-2}, \dots, U^2, X^2, U^1, X^1) \\
&= u * \psi(X^n, U^{n-1}, X^{n-1}, U^{n-2}, X^{n-2}, \dots, U^2, X^2, U^1, X^1) \\
&= u *_c \psi(X^n, U^{n-1}, X^{n-1}, U^{n-2}, X^{n-2}, \dots, U^2, X^2, U^1, X^1) \\
&= Q_u \psi(X^n, U^{n-1}, X^{n-1}, U^{n-2}, X^{n-2}, \dots, U^2, X^2, U^1, X^1)
\end{aligned}$$

for every $(X^n, U^{n-1}, X^{n-1}, U^{n-2}, X^{n-2}, \dots, U^2, X^2, U^1, X^1) \in K_1$, where $*$ signifies the convolution product on N with respect the variables $(X^n, X^{n-1}, X^{n-2}, \dots, X^2, X^1)$ and $*_c$ signifies the commutative convolution product on M with respect the variables $(X^n, U^{n-1}, U^{n-2}, \dots, U^2, U^1)$.

Proof: In fact we have. By Lemma 3.1.1, the mapping $\psi \mapsto Q_u \psi$ is a topological isomorphism of $\mathcal{S}_E^I(K_1)$ onto its image, then the mapping $\psi \mapsto P_u \psi$ is a topological isomorphism of $\mathcal{S}_E^I(K_1)$ onto its image.

Since

$$\begin{aligned}
& R(P_u \psi)(X^n, U^{n-1}, X^{n-1}, U^{n-2}, X^{n-2}, \dots, U^2, X^2, U^1, X^1) \\
&= P_u(R\psi)(X^n, U^{n-1}, X^{n-1}, U^{n-2}, X^{n-2}, \dots, U^2, X^2, U^1, X^1) \quad (18)
\end{aligned}$$

so the following diagram is commutative:

$$\begin{array}{ccc}
\mathcal{S}_E^I(K_1) & \xrightarrow{P_u} & P_u \mathcal{S}_E^I(K_1) \\
\sim \uparrow \downarrow R & & \downarrow R \\
\mathcal{S}(N) & \xrightarrow{Q_u} & P_u \mathcal{S}(N)
\end{array}$$

Hence the mapping $\psi \mapsto P_u \psi$ is a topological isomorphism of $\mathcal{S}(N)$ onto its image. So the transpose ${}^t P_u$ of P_u is a continuous mapping of $\mathcal{S}'(N)$ onto $\mathcal{S}'(N)$. This means that for every tempered distribution T on N there is a tempered distribution E on N such that

$$P_u F = T \quad (19)$$

Indeed the Dirac measure δ belongs to $\mathcal{S}'(N)$.

If we use Atiyah method [1], and [9], we can obtain the following result
"Every invariant differential operator on N which is not identically 0 has a tempered fundamental solution".

4 Fourier Analysis on AN .

4.1. Let $G = SL(n, \mathbb{R})$ be the real semi-simple Lie group and let $G = KAN$ be the Iwasawa decomposition of G , where $K = SO(n, \mathbb{R})$, and

$$N = \begin{pmatrix} 1 & * & . & . & * \\ 0 & 1 & * & . & * \\ . & . & . & . & * \\ . & . & . & . & . \\ 0 & 0 & . & 0 & 1 \end{pmatrix}, \quad (20)$$

$$A = \begin{pmatrix} a_1 & 0 & 0 & . & 0 \\ 0 & a_2 & 0 & . & 0 \\ . & . & . & . & . \\ . & . & . & . & . \\ 0 & 0 & . & 0 & a_n \end{pmatrix} \quad (21)$$

where $a_1.a_2....a_n = 1$ and $a_i \in \mathbb{R}_+^*$. The product AN is a closed subgroup of G and is isomorphic (algebraically and topologically) to the semi-direct product of A and N with N normal in AN .

Then the group AN is nothing but the group $S = N \rtimes_{\rho} A$. So the product of two elements X and Y by

$$(x, a)(m, b) = (x.\rho(a)y, a.b) \quad (22)$$

for any $X = (x, a_1, a_2, .., a_n) \in S$ and $Y = (m, b_1, b_2, .., b_n) \in S$. Let $dnda = dmda_1da_2..da_{n-1}$ be the right haar measure on S and let $L^2(S)$ be the Hilbert space of the group S . Let $L^1(S)$ be the Banach algebra that consists of all complex valued functions on the group S , which are integrable with respect to the Haar measure of S and multiplication is defined by convolution on S as

$$g * f = \int_S f((m, b)^{-1}(n, a))g(m, b)dmdb \quad (23)$$

where $dmdb = dmdb_1db_2..db_{n-1}$ is the right Haar measure on $S = N \rtimes_{\rho} A$.

In the following we prove the Plancherel theorem. Therefore let $T = N \times A$ be the Lie group of direct product of the two Lie groups N and A ,

and let $H = N \times A \times A$ the Lie group, with law

$$(n, t, r)(m, s, q) = (n\rho(r)m, ts, rq) \quad (24)$$

for all $(n, t, r) \in H$ and $(m, s, q) \in H$. In this case the group S can be identified with the closed subgroup $N \times \{0\} \times_\rho A$ of H and T with the subgroup $N \times A \times \{0\}$ of H

Definition 4.1. For every function f defined on S , one can define a function $\tilde{f}$ on L as follows:

$$\tilde{f}(n, a, b) = f(\rho(a)n, ab) \quad (25)$$

for all $(n, a, b) \in H$. So every function $\psi(n, a)$ on S extends uniquely as an invariant function $\tilde{\psi}(n, b, a)$ on L .

Remark 4.1. The function $\tilde{f}$ is invariant in the following sense:

$$\tilde{f}(\rho(s)n, as^{-1}, bs) = \tilde{f}(n, a, b) \quad (26)$$

for any $(n, a, b) \in H$ and $s \in H$.

Lemma 4.1. For every function $f \in L^1(S)$ and for every $g \in L^1(S)$, we have

$$g * \tilde{f}(n, a, b) = g *_c \tilde{f}(n, a, b) \quad (27)$$

$$\int_{\mathbb{R}^{n-1}} \mathcal{F}^d(g * \tilde{f})(\lambda, \mu, \nu) d\nu = \mathcal{F}^d \tilde{f}(\lambda, \mu, 0) \mathcal{F}_A^{n-1} g(\lambda, \mu) \quad (28)$$

for every $(n, a, b) \in H$, where $*$ signifies the convolution product on S with respect the variables (n, b) and $*_c$ signifies the commutative convolution product on B with respect the variables (n, a) , where $\mathcal{F}_A^{n-1}$ is the Fourier transform on A and $\mathcal{F}^d$ is the Fourier transform on N

Proof: In fact we have

$$\begin{aligned} g * \tilde{f}(n, a, b) &= \int_G \tilde{f}((m, c)^{-1}(n, a, b)) g(m, s) dm dc \\ &= \int_S \tilde{f}[(\rho(c^{-1})(m^{-1}), c^{-1})(n, a, b)] u(m, s) dm dc \\ &= \int_S \tilde{f}[\rho(c^{-1})(m^{-1}n), a, bc^{-1}] g(m, c) dm dc \\ &= \int_S \tilde{f}[m^{-1}n, ac^{-1}, b] g(m, c) dm dc = g *_c \tilde{f}(n, a, b) \end{aligned} \quad (29)$$

and so

$$\begin{aligned}
& \int_{\mathbb{R}^{n-1}} \mathcal{F}^d \mathcal{F}_A^{n-1} (g * \tilde{f})(\lambda, \mu, \nu) d\nu \\
&= \int_{\mathbb{R}^{n-1}} \mathcal{F}^d \mathcal{F}_A^{n-1} (g *_c \tilde{f})(\lambda, \mu, \nu) d\nu \\
&= \mathcal{F}^d \mathcal{F}_A^{n-1} \tilde{f}(\lambda, \mu, 0) \mathcal{F}_A^{n-1} g(\lambda, \mu)
\end{aligned} \tag{30}$$

Plancherel Theorem 4.1. *For any function $\Psi \in L^1(S)$, we have*

$$\int_{\mathbb{R}^d} \int_{\mathbb{R}^{n-1}} |\mathcal{F}^d \mathcal{F}_A^{n-1} \Psi(\lambda, \mu)|^2 d\lambda = \int_{AN} |\Psi(X, a)|^2 dX \tag{31}$$

Proof: Let $\widetilde{\Psi}$ the function defined as

$$\widetilde{\Psi}(n, a, b) = \check{\Psi}(an, ab) = \overline{\Psi[(an, ab)^{-1}]} \tag{32}$$

Then we obtain

$$\begin{aligned}
& \Psi * \widetilde{\Psi}(0, 0, 0) \\
&= \int_{NA} \widetilde{\Psi}((n, a)^{-1}(0, 0, 0)) \Psi((n, a) dnda \\
&= \int_{NA} \widetilde{\Psi}(a^{-1}.n, a^{-1})(0, 0, 0)) \Psi((n, a) dnda = \int_{NA} \widetilde{\Psi}(a^{-1}.n, 0, a^{-1}) \Psi((n, a) dnda \\
&= \int_{NA} \widetilde{\Psi}((-X^1)(X), 0, -X^1) \Psi((n, a) dnda = \int_{NA} \check{\Psi}(a^{-1}.n, a^{-1}) \Psi((n, a) dnda \\
&= \int_{NA} \overline{\check{\Psi}(a^{-1}.n, a)^{-1} \Psi((n, a) dnda = \int_{\mathbb{R}^d} |\Psi(n, a)|^2 dnda \tag{33}
\end{aligned}$$

and by lemma 4.1, we get

$$\begin{aligned}
& \Psi_* \widetilde{\Psi}(0, 0, 0) \\
&= \int_N \int_A \mathcal{F}^d \mathcal{F}_A^{n-1}(\Psi_* \widetilde{\Psi})(\xi, \lambda, \mu) d\xi d\lambda d\mu \\
&= \int_N \int_A \mathcal{F}^d \mathcal{F}_A^{n-1}(\Psi_* \widetilde{\Psi})(\xi, \lambda, \mu) d\xi d\lambda d\mu \\
&= \int_N \int_A \mathcal{F}^d \mathcal{F}_A^{n-1} \widetilde{\Psi}(\xi, \lambda, \mu) \mathcal{F}^d \mathcal{F}_A^{n-1} \Psi(\xi, \lambda) d\xi d\lambda d\mu \\
&= \int_N \int_A \int_N \int_A \int_N \int_A \widetilde{\Psi}(n, a, 0) e^{-i\langle \xi, n \rangle} e^{-i\langle \lambda, a \rangle} \Psi(m, b) e^{-i\langle \xi, m \rangle} e^{-i\langle \lambda, b \rangle} \\
&\quad dndadmdbd\xi d\lambda \\
&= \int_N \int_A \int_N \int_A \int_N \int_A \widetilde{\Psi}(an, a) e^{-i\langle \xi, n \rangle} e^{-i\langle \lambda, a \rangle} \Psi(m, b) e^{-i\langle \xi, m \rangle} e^{-i\langle \lambda, b \rangle} \\
&\quad dndadmdbd\xi d\lambda \\
&= \int_N \int_A \int_N \int_A \int_N \int_A \overline{\Psi(an, a)^{-1}} e^{-i\langle \xi, n \rangle} e^{-i\langle \lambda, a \rangle} \Psi(m, b) e^{-i\langle \xi, m \rangle} e^{-i\langle \lambda, b \rangle} \\
&\quad dndadmdbd\xi d\lambda \\
&= \int_N \int_A \int_N \int_A \int_N \int_A \overline{\Psi(n^{-1}, -a) e^{-i\langle \xi, n \rangle} e^{-i\langle \lambda, a \rangle}} \Psi(m, b) e^{-i\langle \xi, m \rangle} e^{-i\langle \lambda, b \rangle} \\
&\quad dndadmdbd\xi d\lambda \\
&= \int_N \int_A \int_N \int_A \int_N \int_A \overline{\Psi(n, a) e^{-i\langle \xi, n \rangle} e^{-i\langle \lambda, a \rangle}} \Psi(m, b) e^{-i\langle \xi, m \rangle} e^{-i\langle \lambda, b \rangle} \\
&\quad dndadmdbd\xi d\lambda \\
&= \int_N \int_A \mathcal{F}^d \mathcal{F}_A^{n-1}(\Psi)(\xi, \lambda) \mathcal{F}^d \mathcal{F}_A^{n-1}(\Psi)(\xi, \lambda) d\xi d\lambda \\
&= \int_N \int_A |\mathcal{F}^d \mathcal{F}_A^{n-1}(\Psi)(\xi, \lambda)|^2 d\xi d\lambda
\end{aligned}$$

where $\langle \xi, n \rangle = \sum_{i=1}^d \xi_i n_i$, and $\langle \lambda, a \rangle = \sum_{j=1}^{n-1} \lambda_j n_j$.

4.2. Let $\mathcal{S}(N \rtimes A)$ be the Schwartz space of $S = N \rtimes A$ which can be considered as the Schwartz space of $\mathcal{S}(T = N \times A)$, let $\mathcal{S}'(N \rtimes A)$ be the space of all tempered distributions on $N \rtimes A$ and let $\mathcal{S}'(T)$ be the space of all tempered distributions on C . If we consider the group S as a subgroup of H , then $\tilde{f} \in \mathcal{S}(S)$ for a is fixed, and if we consider T as a subgroup of H , then $\tilde{f} \in \mathcal{S}(T)$ for b is fixed. This being so; denote by $\mathcal{S}_E(H)$ the space of all functions $\Phi(x, a, b) \in C^\infty(H)$ such that $\Phi(x, a, b) \in \mathcal{S}(N)$ for a is fixed, and $\Phi(x, a, b) \in \mathcal{S}(T)$ for b is fixed. We equip $\mathcal{S}_E(H)$ with the natural topology defined by the seminormas:

$$\Phi \rightarrow \sup_{(x,a) \in T} |Q(Y)P(D)\Phi(x, a, b)| \quad b \text{ fixed}$$

$$\Phi \rightarrow \sup_{(x,b) \in N} |R(Y)S(D)\Phi(x, a, b)| \quad a \text{ fixed}$$

where P, Q, R and S run over the family of all complex polynomials in $2d-1$ variables. Let $\mathcal{S}_E^I(H)$ be the subspace of all functions $\psi \in \mathcal{S}_E(H)$, which are invariant in sense (13), then we have the following result.

Proposition 4.1. *Let $u \in \mathcal{U}$ and D_u be the invariant differential operator on the group T , which is associated to u , acts on the variables $(x, a) \in T$, then we have*

(i) *The mapping $f \mapsto \tilde{f}$ is a topological isomorphism of $\mathcal{S}(N)$ onto $\mathcal{S}_E^I(H)$.*

(ii) *The mapping $\Phi \mapsto Q_u \Phi$ is a topological isomorphism of $\mathcal{S}_E^I(H)$ onto its image*

Proof: (i) In fact $\sim$ is continuous and the restriction mapping $\Phi \mapsto R\Phi$ on S is continuous from $\mathcal{S}_E^I(H)$ into $\mathcal{S}(S)$ that satisfies $R \circ \sim = Id_{\mathcal{S}(N \rtimes A)}$ and $\sim \circ R = Id_{\mathcal{S}_E^I(H)}$, where $Id_{\mathcal{S}(S)}$ (resp. $Id_{\mathcal{S}_E^I(H)}$) is the identity mapping of $\mathcal{S}(S)$ (resp. $\mathcal{S}_E^I(H)$) and S is considered as a subgroup of H .

To prove(ii) we refer to [16, P.313 – 315] and his famous result that is: "Any invariant differential operator on T , is a topological isomorphism of $\mathcal{S}(T)$ onto its image" From this result, we obtain that

$$D_u : \mathcal{S}_E(H) \rightarrow \mathcal{S}_E(H) \quad (34)$$

is a topological isomorphism and its restriction on $\mathcal{S}_E^I(H)$ is a topological isomorphism of $\mathcal{S}_E^I(H)$ onto its image. Hence the proposition is proved.

In the following we will prove that every invariant differential operator on $S = N \rtimes A$ has a tempered fundamental solution. We consider the two invariant differential operators P_u and Q_u , the first on the group S , and the second on the abelian vector group T . Our main result is:

Existence theorem 4.2. *Every nonzero invariant differential operator on $N \rtimes A$ has a tempered fundamental solution*

Proof: For every function $\psi \in C^\infty(H)$ invariant in sense (11) and for every $u \in \mathcal{U}$, we have

$$\begin{aligned} P_u \psi(x, a, b) &= u * \psi(x, a, b) \\ &= u *_c \psi(x, a, b) = Q_u \psi(x, a, b) \end{aligned}$$

for every $(x, a, b) \in H$, where $*$ signifies the convolution product on S with respect the variables (x, b) and $*_c$ signifies the commutative convolution product on T with respect the variables (x, a) .

Proof: In fact we have by proposition 4.1, the mapping $\psi \mapsto Q_u \psi$ is a topological isomorphism of $\mathcal{S}_E^I(H)$ onto its image, then the mapping $\psi \mapsto P_u \psi$ is a topological isomorphism of $\mathcal{S}_E^I(H)$ onto its image.

Since

$$R(P_u \psi)(x, a, b) = P_u(R\psi)(x, a, b) \quad (35)$$

so the following diagram is commutative:

$$\begin{array}{ccc} \mathcal{S}_E^I(H) & \xrightarrow{P_u} & P_u \mathcal{S}_E^I(H) \\ \sim \uparrow \downarrow R & & \downarrow R \\ \mathcal{S}(N \rtimes A) & \xrightarrow{Q_u} & P_u \mathcal{S}(N \rtimes A) \end{array}$$

Hence the mapping $\psi \mapsto P_u \psi$ is a topological isomorphism of $\mathcal{S}(N \rtimes A)$ onto its image. So the transpose ${}^t P_u$ of P_u is a continuous mapping of $\mathcal{S}'(N \rtimes A)$ onto $\mathcal{S}'(N \rtimes A)$. This means that for every tempered distribution F on $N \rtimes A$ there is a tempered distribution E on $N \rtimes A$ such that

$$P_u E = F \quad (36)$$

Indeed the Dirac measure δ belongs to $\mathcal{S}'(N \rtimes A)$.

5 Ideals Algebra $L^1(N)$.

Lemma 5.1. (i) The mapping Γ from $\widetilde{L^1(N)}|_M$ to $\widetilde{L^1(N)}|_N$ defined by

$$\begin{aligned} & \widetilde{\phi}|_M (X^n, X^{n-1}, 0, X^{n-2}, 0, \dots, X^2, 0, X^1, 0) \\ \rightarrow & \Gamma(\widetilde{\phi}|_M)(X^n, 0, X^{n-1}, 0, X^{n-2}, \dots, 0, X^2, 0, X^1) \\ = & \widetilde{\phi}|_N(X^n, 0, X^{n-1}, 0, X^{n-2}, \dots, 0, X^2, 0, X^1) \end{aligned}$$

is a topological isomorphism

(ii) For every $\psi \in L^1(N)$ and $\phi \in L^1(N)$, we obtain

$$\begin{aligned} & \Gamma(\psi *_c \widetilde{\phi}|_M)(X^n, 0, X^{n-1}, 0, X^{n-2}, \dots, 0, X^2, 0, X^1) \\ = & \psi *_c \widetilde{\phi}|_N(X^n, 0, X^{n-1}, 0, X^{n-2}, \dots, 0, X^2, 0, X^1) \\ = & \psi *_c \widetilde{\phi}(X^n, X^{n-1}, X^{n-2}, \dots, X^2, X^1) \end{aligned}$$

$$\begin{aligned} & (\psi *_c \widetilde{\phi}|_M)(X^n, X^{n-1}, 0, X^{n-2}, 0, \dots, X^2, 0, X^1, 0) \\ = & \int_M \widetilde{\phi} [(X^n - Y^n, X^{n-1} - Y^{n-1}, 0, X^{n-2} - Y^{n-2}, 0, \dots, X^2 - Y^2, 0, X^1 - Y^1, 0] \\ & \psi(Y^n, Y^{n-1}, Y^{n-2}, \dots, Y^2, Y^1) dY^n dY^{n-1} dY^{n-2}, \dots, dY^2 dY^1 \end{aligned}$$

Proof: (i) The mapping Γ is continuous and has an inverse Γ^{-1} given by

$$\begin{aligned} & \widetilde{\phi}|_N (X^n, 0, X^{n-1}, 0, X^{n-2}, \dots, 0, X^2, 0, X^1) \\ \rightarrow & \Gamma^{-1}(\widetilde{\phi}|_N)(X^n, X^{n-1}, 0, X^{n-2}, 0, \dots, X^2, 0, X^1, 0) \\ = & \widetilde{\phi}|_M(X^n, X^{n-1}, 0, X^{n-2}, 0, \dots, X^2, 0, X^1, 0) \end{aligned}$$

(ii) It is enough to see for every $\phi \in L^1(N)$

$$\begin{aligned} & \Gamma(\psi *_c \widetilde{\phi}|_M)(X^n, 0, X^{n-1}, 0, X^{n-2}, \dots, 0, X^2, 0, X^1) \\ = & \int_M \widetilde{\phi}|_M [X^n - Y^n, -Y^{n-1}, X^{n-1}, -Y^{n-2}, X^{n-2}, \dots, -Y^2, X^2, -Y^1, X^1] \\ & \psi(Y^n, Y^{n-1}, Y^{n-2}, \dots, Y^2, Y^1) dY^n dY^{n-1} dY^{n-2}, \dots, dY^2 dY^1 \\ = & \psi *_c \phi(X^n, X^{n-1}, X^{n-2}, \dots, X^2, X^1), \quad \phi \in L^1(N) \end{aligned}$$

If I is a subalgebra of $L^1(N)$, we denote by $\tilde{I}$ its image by the mapping $\sim$. Let $J = \tilde{I}|_M$. Our main result is:

Theorem 5.1. *Let I be a subalgebra of $L^1(N)$, then the following conditions are equivalents.*

(i) $J = \tilde{I}|_M$ is an ideal in the Banach algebra $L^1(M)$.

(ii) I is a left ideal in the Banach algebra $L^1(N)$.

Proof: (i) implies (ii) Let I be a subspace of the space $L^1(N)$ such that $J = \tilde{I}|_M$ is an ideal in $L^1(M)$, then we have:

$$\begin{aligned} & \psi *_c \tilde{I}|_M(X^n, X^{n-1}, 0, X^{n-2}, 0, \dots, X^2, 0, X^1, 0) \\ & \subseteq \tilde{I}|_M(X^n, X^{n-1}, 0, X^{n-2}, 0, \dots, X^2, 0, X^1, 0) \end{aligned} \quad (37)$$

for any $\psi \in L^1(M)$ and $(x, x_3, x_2, x_1) \in M$, where

$$\begin{aligned} & \psi *_c \tilde{I}|_M(X^n, X^{n-1}, 0, X^{n-2}, 0, \dots, X^2, 0, X^1, 0) \\ = & \left\{ \begin{array}{l} \int_M \tilde{\phi}[X^n - Y^n, -Y^{n-1}, X^{n-1}, -Y^{n-2}, X^{n-2}, \dots, -Y^2, X^2, -Y^1, X^1] \\ \psi(Y^n, Y^{n-1}, Y^{n-2}, \dots, Y^2, Y^1 \\ dY^n dY^{n-1} dY^{n-2}, \dots, dY^2 dY^1, \quad \phi \in L^1(N) \end{array} \right\} \end{aligned}$$

It shows that

$$\begin{aligned} & \psi *_c \tilde{\phi}|_M(X^n, X^{n-1}, 0, X^{n-2}, 0, \dots, X^2, 0, X^1, 0) \\ & \in \tilde{I}|_M(X^n, X^{n-1}, 0, X^{n-2}, 0, \dots, X^2, 0, X^1, 0) \end{aligned} \quad (38)$$

for any $\tilde{\phi} \in \tilde{I}$. Then we get

$$\begin{aligned} & \Gamma(\psi *_c \tilde{\phi}|_M)(X^n, 0, X^{n-1}, 0, X^{n-2}, \dots, 0, X^2, 0, X^1) \\ = & \psi *_c \tilde{\phi}(X^n, 0, X^{n-1}, 0, X^{n-2}, \dots, 0, X^2, 0, X^1) \\ \in & \Gamma(\tilde{I}|_M)(X^n, 0, X^{n-1}, 0, X^{n-2}, \dots, 0, X^2, 0, X^1) \\ = & \tilde{I}|_N(X^n, 0, X^{n-1}, 0, X^{n-2}, \dots, 0, X^2, 0, X^1) \\ = & I(X^n, X^{n-1}, X^{n-2}, \dots, X^2, X^1) \end{aligned} \quad (39)$$

It is clear that (ii) implies (i).

Corollary 5.1. *Let I be a subset of the space $L^1(N)$ and $\tilde{I}$ its image by the mapping $\sim$ such that $J = \tilde{I}|_M$ is an ideal in $L^1(N)$, then the following conditions are verified.*

(i) J is a closed ideal in the algebra $L^1(M)$ if and only if I is a left closed ideal in the algebra $L^1(N)$.

(ii) J is a maximal ideal in the algebra $L^1(M)$ if and only if I is a left maximal ideal in the algebra $L^1(N)$.

(iii) J is a prime ideal in the algebra $L^1(M)$ if and only if I is a left prime ideal in the algebra $L^1(N)$.

(iv) J is a dense ideal in the algebra $L^1(M)$ if and only if I is a left dense ideal in the algebra $L^1(N)$.

The proof of this corollary results immediately from theorem 5.1.

5.2. Some New Groups

I- Let $\mathbb{R}_-^* = \{x \in \mathbb{R}; x \neq 0\}$, with law

$$x \cdot y = -x \cdot y \quad (40)$$

for all $x \in \mathbb{R}_-^*$ and $y \in \mathbb{R}_-^*$, where $\cdot$ signifies the product in $\mathbb{R}_-^*$ and $\cdot$ signifies the ordinary product of two real numbers.

$(\mathbb{R}_-^*, \cdot)$ with this new law $\cdot$ becomes a commutative group isomorphic with the multiplicative group $(\mathbb{R}_+^*, \cdot)$ and with the real vector group $(\mathbb{R}, +)$.

Proof: the identity element is -1 because

$$\begin{aligned} (-1) \cdot x &= -(-1) \cdot x = x, \text{ and} \\ x \cdot (-1) &= -x \cdot (-1) = x \end{aligned} \quad (41)$$

If $(x, y) \in \mathbb{R}_-^* \times \mathbb{R}_-^*$, such that $x \cdot y = -1$, then we get

$$x \cdot y = -x \cdot y = -1 \quad (42)$$

From this equation we obtain $y = \frac{1}{x} = x^{-1} \in \mathbb{R}_-^*$, which is the inverse of x . Now let x, y , and z be three elements belong to $\mathbb{R}_-^*$, then we have

$$(x \cdot y) \cdot z = -(x \cdot y) \cdot z = -(-x \cdot y) \cdot z = x \cdot y \cdot z \quad (43)$$

and

$$x \cdot (y \cdot z) = x \cdot (-y \cdot z) = -x \cdot (-y \cdot z) = x \cdot y \cdot z \quad (44)$$

So the law is associative.

Clearly the law $\cdot$ is commutative because

$$x \cdot y = -x \cdot y = -y \cdot x = y \cdot x \quad (45)$$

Let $\psi : \mathbb{R}_-^* \rightarrow \mathbb{R}_+^*$ the mapping defined by

$$\psi(x) = -x \quad (46)$$

then we get

$$\begin{aligned} \psi(x \cdot y) &= -(x \cdot y) \\ &= x \cdot y = (-x) \cdot (-y) = \psi(x) \cdot \psi(y) \end{aligned} \quad (47)$$

Another hand we have

$$\Psi(x) = \Psi(y) \implies -x = -y \implies x = y \quad (48)$$

and if $y \in \mathbb{R}_+^*$, there is $x \in \mathbb{R}_-^*$, such that $\Psi(x) = y$. In fact it is enough to take $x = -y$

That means ψ is a group isomorphism from $\mathbb{R}_-^*$ onto $\mathbb{R}_+^*$. Since the group $(\mathbb{R}_+^*, \cdot)$ is isomorphic with the real vector group $(\mathbb{R}, +)$, so the new group $\mathbb{R}_-^*$ is isomorphic with the group $(\mathbb{R}, +)$

Then $\mathbb{R}^* = \mathbb{R}_-^* \cup \mathbb{R}_+^* = \mathbb{R}_+^* \cup \mathbb{R}_+^*$

II- The group $\mathbb{R}_+^* \times \mathbb{R}_-^*$, which is the direct product of $\mathbb{R}_+^*$ and $\mathbb{R}_-^*$ is topological and algebraic isomorphic onto $(\mathbb{R}^2, +)$

Proof: Let $\Psi : \mathbb{R}_+^* \times \mathbb{R}_-^* \rightarrow \mathbb{R}^2$ the mapping defined by

$$\Psi(x, y) = (\ln x, \ln |y|) \quad (49)$$

The mapping Ψ is one-to-one and onto. Now, let us to prove that Ψ is group homomorphism

$$\begin{aligned} \Psi((x, y)(s, t)) &= \Psi((x.s, y \bullet t) = (\ln(x.s), \ln |y \bullet t|) \\ &= (\ln(x.s), \ln |(-1)y.t|) = (\ln x + \ln s, \ln |y.t|) \\ &= (\ln x + \ln s, \ln |y| + \ln |t|) = (\ln x, \ln |y|) + (\ln s, \ln |t|) \\ &= \Psi((x, y) + \Psi(s, t)) \end{aligned}$$

The inverse of Ψ is given by

$$\Psi^{-1}(a, b) = (e^a, -e^b) \quad (50)$$

III- The group $\mathbb{R}_+^* \times \mathbb{R}_-^*$ is also isomorphic onto the group $(\mathbb{C}, +)$

Proof: It is enough to consider the mapping $\Phi : \mathbb{R}_+^* \times \mathbb{R}_-^* \rightarrow \mathbb{C}$ defined by $\Phi(x, y) = \ln x + \ln |y|^i$, $i = (-1)^{\frac{1}{2}}$.

Acknowledgment. For all the results of this paper the authors advise the readers to refer my paper [9],

References

- [1] M. F. Atiyah., (1970), *Resolution of Singularities and Division of Distributions*, Comm, on Pure and App. Math, Vol. 23. 145-150.
- [2] K. El- Hussein., (2009), *A Fundamental Solution of an Invariant Differential Operator on the Heisenberg Group*, International Mathematical Forum, 4, no. 12, 601- 612.
- [3] K. El- Hussein., (2011), *On the Left Ideals of Group Algebra of thje Affine Group*. International Mathematical Forum, Vol 6. no. 4, 193-204.
- [4] K. El- Hussein., (1989), *Opérateurs Différentiels Invariants sur les Groupes de Deplacements*, Bull. Sc. Math. 2^e series 113, 89-117.
- [5] K. El- Hussein., (2009), *Eigendistributions for the Invariant Differential Operators on the Affine Group*, Int. Journal of Math. Analysis, Vol. 3, no. 9 , 419-429.
- [6] K. El- Hussein, (2010), *Note on the Solvability of Lewy Operator*, Int. Math. Forum, 5, no. 8, 389 - 393.
- [7] K. El- Hussein., (2009), *Eigendistributions for the Invariant Differential operators on the Affine Group*. Int. Journal of Math. Analysis, Vol. 3, no. 9, 419-429.
- [8] K. El- Hussein., (2010), *Fourier transform and invariant differential operators on the solvable Lie group G_4* , in Int. J. Contemp. Maths Sci. 5. No. 5-8, 403-417.
- [9] K. El- Hussein., (2013), *Abstract Harmonic Analysis and Ideals of Banach Algebra on 3-Step Nilpotent Lie Group*, International Journal of Engineering Research & Technology (IJERT), Vol. 2 Issue 11, November.
- [10] K. El- Hussein., (2014), *Abstract Harmonic Analysis on Spacetime*, arXiv:submit/0951116 [math-ph] 6 Apr
- [11] Harish-Chandra; (1952), *Plancherel formula for 2×2 real unimodular group*, Proc. nat. Acad. Sci. U.S.A., vol. 38, pp. 337-342.

- [12] Harish-Chandra; (1952), *The Plancherel formula for complex semi-simple Lie group*, Trans. Amer. Mth. Soc., vol. 76, pp. 485- 528.
- [13] S. Helgason., 1984, *Groups and Geometric Analysis*, Academic Press.
- [14] S. Helgason., (2005), *The Abel, Fourier and Radon Transforms on Symmetric Spaces*. Indagationes Mathematicae. 16, 531-551.
- [15] L. Hormander, 1983, *The analysis of Linear Partial Differential Operator I*”, Springer-Verlag.
- [16] F. Trèves., (1962), *Linear Partial Differential Equations with Constant Coefficients*, Gordon and Breach.
- [17] W. Rudin., (1962), *Fourier Analysis on Groups*, Interscience Publishers, New York, NY.G.
- [18] Warner., (1970), *Harmonic Analysis on Semi-Simple Lie Groups*, Springer-verlag Berlin heidelberg New york.